arXiv:0809.1889v1 [stat.ME] 10 Sep 2008

The Beta Generalized Exponential Distribution

Wagner Barreto-Souza[a], Alessandro H. S. Santos[b] and Gauss M. Cordeiro[b]

[a] *Departamento de Estatística*,
Universidade Federal de Pernambuco,
Cidade Universitária, 50740-540 – Recife, PE, Brazil
(wagnerbs85@hotmail.com);

[b] *Departamento de Estatística e Informática*,
Universidade Federal Rural de Pernambuco,
Rua Dom Manoel de Medeiros s/n, 50171-900 – Recife, PE, Brazil
(alessandrohss@yahoo.com.br, gausscordeiro@uol.com.br)

Abstract

We introduce the beta generalized exponential distribution that includes the beta exponential and generalized exponential distributions as special cases. We provide a comprehensive mathematical treatment of this distribution. We derive the moment generating function and the rth moment thus generalizing some results in the literature. Expressions for the density, moment generating function and rth moment of the order statistics also are obtained. We discuss estimation of the parameters by maximum likelihood and provide the information matrix. We observe in one application to real data set that this model is quite flexible and can be used quite effectively in analyzing positive data in place of the beta exponential and generalized exponential distributions.

keywords: Beta exponential distribution, Information matrix, Generalized exponential distribution, Maximum likelihood estimation.

1 Introduction

Gupta and Kundu [4] defined the cumulative distribution function (cdf) of the generalized exponential (GE) distribution by

$$G_{\lambda,\alpha}(x) = (1 - e^{-\lambda x})^{\alpha}, \quad x > 0. \tag{1}$$

The two parameters of the GE distribution represent the shape ($\alpha > 0$) and the scale parameter ($\lambda > 0$) like the gamma and Weibull distributions. The distribution (1) is also named the exponentiated exponential distribution. Clearly, the exponential distribution

is a particular case of the GE distribution when $\alpha = 1$. Gupta and Kundu also investigated some of their properties. The exponentiated Weibull (EW) distribution introduced by Mudholkar et al. [7] extends the GE distribution and was also studied by Mudholkar et al. [8], Mudholkar and Hutson [9] and Nassar and Eissa [13]. Nadarajah and Kotz [12] introduced four more exponentiated type distributions: the exponentiated gamma, exponentiated Weibull, exponentiated Gumbel and exponentiated Fréchet distributions by generalizing the gamma, Weibull, Gumbel and Fréchet distributions in the same way that the GE distribution extends the exponential distribution. They also provide some mathematical properties for each exponentiated distribution.

The GE density function varies significantly depending on the shape parameter α. Also, the hazard function is a non-decreasing function if $\alpha > 1$, and it is a non-increasing function if $\alpha < 1$. For $\alpha = 1$, it is constant. The GE distribution has lots of properties which are quite similar in nature to those of the gamma distribution but it has explicit expressions for the distribution and survival functions like a Weibull distribution. The gamma, Weibull and GE distributions extend the exponential distribution but in different ways. Therefore, it can be used as an alternative to the Weibull and gamma distributions and in some situations it might work better in terms of fitting than the other two distributions although it can not be guaranteed. Moreover, it is well known that the gamma distribution has certain advantages compared to the Weibull in terms of the faster convergence of the maximum likelihood estimates (MLEs). It is expected that the GE distribution also should enjoy those properties.

Consider starting from the cdf $G(x)$ of a random variable, Eugene et al. [2] defined a class of generalized distributions from it given by

$$F(x) = \frac{1}{B(a,b)} \int_0^{G(x)} \omega^{a-1}(1-\omega)^{b-1} d\omega, \tag{2}$$

where $a > 0$ and $b > 0$ are two additional parameters whose role is to introduce skewness and to vary tail weight and $B(a,b) = \int_0^1 \omega^{a-1}(1-\omega)^{b-1} d\omega$ is the beta function. The cdf $G(x)$ could be quite arbitrary and F is named the beta G distribution. Application of $X = G^{-1}(V)$ to $V \sim B(a,b)$ yields X with cdf (2).

Eugene et al. [2] defined the beta normal (BN) distribution by taking $G(x)$ in (2) to be the cdf of the normal distribution and derived some first moments. General expressions for the moments of the BN distribution were derived by Gupta and Nadarajah [6]. Nadarajah and Kotz [10] considered the beta Gumbel (BG) distribution by taking $G(x)$ as the cdf of the Gumbel distribution and provided closed-form expressions for the moments, the asymptotic distribution of the extreme order statistics and discussed the maximum likelihood estimation procedure. Also, Nadarajah and Kotz [11] worked with the beta exponential (BE) distribution and obtained the moment generating function, the first four cumulants, the asymptotic distribution of the extreme order statistics and discussed the maximum likelihood estimation. Some of the Nadarajah and Kotz's [11] results were generalized by Cordeiro et al. [1] who considered the beta Weibull distribution. They derived the moment generating function, the moments and the information matrix.

We can write (2) by

$$F(x) = I_{G(x)}(a,b), \tag{3}$$

where $I_y(a,b) = B(a,b)^{-1}\int_0^y w^{a-1}(1-w)^{b-1}dw$ denotes the incomplete beta function ratio, i.e., the cdf of the beta distribution with parameters a and b. For general a and b, we can express (3) in terms of the well-known hypergeometric function defined by

$$ {}_2F_1(\alpha,\beta,\gamma;x) = \sum_{i=0}^{\infty} \frac{(\alpha)_i(\beta)_i}{(\gamma)_i i!}\, x^i, $$

where $(\alpha)_i = \alpha(\alpha+1)\ldots(\alpha+i-1)$ denotes the ascending factorial. We obtain

$$ F(x) = \frac{G(x)}{a\,B(a,b)}\, {}_2F_1(a,1-b,a+1;G(x)). $$

The properties of $F(x)$ for any beta G distribution defined from a parent $G(x)$ in (2) could, in principle, follow from the properties of the hypergeometric function which are well established in the literature; see, for example, Section 9.1 of Gradshteyn and Ryzhik [3].

The probability density function (pdf) corresponding to (2) can be put in the form

$$ f(x) = \frac{1}{B(a,b)} G(x)^{a-1}\{1-G(x)\}^{b-1} g(x), \tag{4} $$

where we noted that $f(x)$ will be most tractable when the cdf $G(x)$ and the pdf $g(x) = dG(x)/dx$ have simple analytic expressions. Except for some special choices for $G(x)$ in (2) as is the case when $G(x)$ is given by (1), it would appear that the pdf $f(x)$ will be difficult to deal with.

We now introduce the four parameter beta generalized exponential (BGE) distribution by taking $G(x)$ in (2) to be the cdf (1). The cdf of the BGE distribution is then

$$ F(x) = \frac{1}{B(a,b)} \int_0^{(1-e^{-\lambda x})^{\alpha}} \omega^{a-1}(1-\omega)^{b-1}d\omega, \quad x>0, \tag{5} $$

for $a>0$, $b>0$, $\lambda>0$ and $\alpha>0$. The pdf and the hazard rate function of the new distribution are, respectively,

$$ f(x) = \frac{\alpha\lambda}{B(a,b)} e^{-\lambda x}(1-e^{-\lambda x})^{\alpha a-1}\{1-(1-e^{-\lambda x})^{\alpha}\}^{b-1}, \quad x>0, \tag{6} $$

and

$$ h(x) = \frac{\alpha\lambda e^{-\lambda x}(1-e^{-\lambda x})^{\alpha a-1}\left\{1-(1-e^{-\lambda x})^{\alpha}\right\}^{b-1}}{B(a,b)\, I_{1-(1-e^{-\lambda x})^{\alpha}}(a,b)}, \quad x>0. \tag{7} $$

The density function (6) does not involve any complicated function. If X is a random variable with pdf (6), we write $X \sim BGE(a,b,\lambda,\alpha)$. The BGE distribution generalizes some well-known distributions in the literature. The GE distribution is a special case for the choice $a=b=1$. If in addition $\alpha=1$, we obtain the exponential distribution with parameter λ. The BE distribution is obtained from (5) with $\alpha=1$. It is evident that (6) is much more flexible than the GE and BE distributions. Plots of the density (6) and failure rate (7) for some special values of a, b, λ and α are given in Figures 1 and 2, respectively. A characteristic of the BGE distribution is that its failure rate



Figure 1: Plots of the density (6) for some values of the parameters.

can be bathtub shaped, monotonically increasing or decreasing and upside-down bathtub depending basically on the values of its parameters.

We now introduce the double generalized exponential (DGE) distribution for the choice $a = 1$ in (5). The cdf of the DGE distribution is

$$F(x) = \{1 - (1 - e^{-\lambda x})^{\alpha}\}^{b}, \quad x > 0. \tag{8}$$

The rest of the paper is organized as follows. In Section 2, we derive some expressions for the cdf of the BGE distribution and for the pdfs of the order statistics. In Sections 3 and 4 we provide expressions for the moment generating function and for the rth moment, respectively. Furthermore, in these sections, we derive corresponding expressions for the order statistics. In Section 5, we discuss maximum likelihood estimation, inference on the parameters, and calculate the information matrix. One application to real data set is presented in Section 6. Some conclusions are drawn in Section 7.



Figure 2: Plots of the hazard (7) for some values of the parameters.

2 Distribution function and order statistics

We provide two simple formulae for the cdf of the BGE distribution depending if the parameter $b > 0$ is real non-integer or integer. First, if $|z| < 1$ and $b > 0$ is real non-integer, we have

$$(1-z)^{b-1} = \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(b)}{\Gamma(b-j)j!} z^j. \tag{9}$$

Using the expansion (9) in (5), the cdf of the BGE distribution when $b > 0$ is real non-integer follows

$$\begin{aligned} F(x) &= \frac{1}{B(a,b)} \int_0^{(1-e^{-\lambda x})^{\alpha}} \omega^{a-1}(1-\omega)^{b-1} d\omega \\ &= \frac{\Gamma(b)}{B(a,b)} \sum_{j=0}^{\infty} \frac{(-1)^j}{\Gamma(b-j)j!} \int_0^{(1-e^{-\lambda x})^{\alpha}} \omega^{a+j-1} d\omega \end{aligned}$$

and then

$$F(x) = \frac{\Gamma(a+b)}{\Gamma(a)} \sum_{j=0}^{\infty} \frac{(-1)^j (1-e^{-\lambda x})^{\alpha(a+j)}}{\Gamma(b-j)j!(a+j)}. \tag{10}$$

Equation (10) reveals the property that the cdf of the BGE distribution can be expressed as an infinite weighted sum of cdfs of GE distributions

$$F(x) = \frac{\Gamma(a+b)}{\Gamma(a)} \sum_{j=0}^{\infty} \frac{(-1)^j G_{\lambda,\alpha(a+j)}(x)}{\Gamma(b-j)j!(a+j)}.$$

Using the binomial expansion in (5), we have when $b > 0$ is integer

$$F(x) = \frac{1}{B(a,b)} \sum_{j=0}^{b-1} \binom{b-1}{j} \frac{(-1)^j}{a+j} (1-e^{-\lambda x})^{\alpha(a+j)}, \tag{11}$$

and, again, the same property of equation (10) holds but now the sum is finite. Expressions (10) and (11) are the main results of this section. The cdf of the BE distribution follows with $\alpha = 1$ from (10) and (11) depending if $b > 0$ is real non-integer and integer, respectively. The cdf of the DGE distribution follows from the above expressions when $a = 1$. The cdf (1) of the GE distribution comes from (11) when $a = b = 1$. If, in addition, $\alpha = 1$, (11) yields the cdf $F(x) = 1 - e^{-\lambda x}$ of the exponential distribution.

It can be seen in the Wolfram Functions Site[1] that for integer a

$$I_y(a,b) = 1 - \frac{(1-y)^b}{\Gamma(b)} \sum_{j=0}^{a-1} \frac{\Gamma(b+j)}{j!} y^j,$$

and for integer b

$$I_y(a,b) = \frac{y^a}{\Gamma(a)} \sum_{j=0}^{b-1} \frac{\Gamma(a+j)}{j!} (1-y)^j.$$

[1] `http://functions.wolfram.com/`

Therefore, for $a > 0$ integer,

$$F(x) = 1 - \frac{\left\{1-(1-e^{-\lambda x})^{\alpha}\right\}^{b}}{\Gamma(b)} \sum_{j=0}^{a-1} \frac{\Gamma(b+j)}{j!}(1-e^{-\lambda x})^{\alpha j}$$

and for $b > 0$ integer we have an alternative form for (11) given by

$$F(x) = \frac{(1-e^{-\lambda x})^{a\alpha}}{\Gamma(a)} \sum_{j=0}^{b-1} \frac{\Gamma(a+j)}{j!}\{1-(1-e^{-\lambda x})^{\alpha}\}^{j}.$$

The density function (6) can be expressed in the mixture form in terms of cdfs of the GE distributions

$$f(x) = \frac{\alpha\lambda}{B(a,b)} e^{-\lambda x} G_{\lambda,a\alpha-1}(x) \sum_{j=0}^{\infty} \frac{(-1)^{j}\Gamma(b)}{\Gamma(b-j)j!} G_{\lambda,\alpha j}(x)$$

for b real non-integer and

$$f(x) = \frac{\alpha\lambda}{B(a,b)} e^{-\lambda x} G_{\lambda,a\alpha-1}(x) \sum_{j=0}^{b-1} \binom{b-1}{j} (-1)^{j} G_{\lambda,\alpha j}(x)$$

for b integer.

Simulation of the BGE distribution is easy: if V is a random variable with a beta distribution with parameters a and b, then $X = -\log(1-V^{1/\alpha})/\lambda$ follows the BGE distribution with parameters a, b, λ and α.

We now give the density of the ith order statistic $X_{i:n}$, $f_{i:n}(x)$ say, in a random sample of size n from the BGE distribution. It is well known that

$$f_{i:n}(x) = \frac{1}{B(i,n-i+1)} f(x) F^{i-1}(x) \left\{1-F(x)\right\}^{n-i},$$

for $i = 1, \ldots, n$. Using (3) and (6) we can express $f_{i:n}(x)$ in terms of the hypergeometric functions by

$$\begin{aligned} f_{i:n}(x) &= \frac{\alpha\lambda e^{-\lambda x} G_{\lambda,a\alpha-1}(x) G_{\lambda,\alpha a(i-1)}(x) \{1-G_{\lambda,\alpha}(x)\}^{b(n-i+1)-1}}{B(i,n-i+1)B(a,b)^{n} a^{i-1} b^{n-i}} \times \\ &\quad {}_2F_1(a, 1-b, a+1; G_{\lambda,\alpha}(x))^{i-1}\, {}_2F_1(b, 1-a, b+1; 1-G_{\lambda,\alpha}(x))^{n-i}. \end{aligned}$$

We now derive two alternative expressions for the densities of the order statistics using the expansion $(\sum_{i=1}^{\infty} a_i)^k = \sum_{\{m_1,\ldots,m_k\}=0}^{\infty} a_{m_1} \ldots a_{m_k}$ for k a positive integer. With this expansion and from (10)−(11), we can show for $b > 0$ real non-integer and integer that

$$f_{i:n}(x) = \sum_{k=0}^{n-i} \sum_{m_1=0}^{\infty} \cdots \sum_{m_{k+i-1}=0}^{\infty} \delta_{k,i}^{(1)} f_{k,i}(x) \tag{12}$$

and

$$f_{i:n}(x) = \sum_{k=0}^{n-i} \sum_{m_1=0}^{b-1} \cdots \sum_{m_{k+i-1}=0}^{b-1} \delta_{k,i}^{(2)} f_{k,i}(x), \tag{13}$$

respectively, where from now on $f_{k,i}(x)$ represents the density of a random variable $X_{k,i}$ following a $BGE(\alpha\{a(i+1) + \sum_{j=1}^{k+i-1} m_j\}, b, \lambda, \alpha)$ distribution, and the functions $\delta_{k,i}^{(1)}$ and $\delta_{k,i}^{(2)}$ required for the above expressions are

$$\delta_{k,i}^{(1)} = \frac{(-1)^{k+\sum_{j=1}^{k+i-1} m_j} \binom{n-i}{k} B(\alpha\{a(i+1) + \sum_{j=1}^{k+i-1} m_j\}, b) \Gamma(b)^{k+i-1}}{B(a,b)^{k+i} B(i, n-i+1) \prod_{j=1}^{k+i-1} \Gamma(b-m_j) m_j! (a+m_j)}$$

and

$$\delta_{k,i}^{(2)} = \frac{(-1)^{k+\sum_{j=1}^{k+i-1} m_j} \binom{n-i}{k} B(\alpha\{a(i+1) + \sum_{j=1}^{k+i-1} m_j\}, b)}{B(a,b)^{k+i} B(i, n-i+1)} \prod_{j=1}^{k+j-1} \frac{\binom{b-1}{m_j}}{(a+m_j)}.$$

The sums in (12) and (13) extends over all $(k+i)$-tuples $(k, m_1, \ldots, m_{k+i-1})$ of non-negative integers and are easily implementable on a computer.

3 Moment generating function

The moment generating function (mgf) of the BGE distribution is given by

$$M(t) = \frac{\alpha\lambda}{B(a,b)} \int_0^\infty e^{tx} e^{-\lambda x} (1 - e^{-\lambda x})^{\alpha a - 1} \{1 - (1 - e^{-\lambda x})^\alpha\}^{b-1} dx. \tag{14}$$

Using the expansion (9) when $b > 0$ is real non-integer, (14) reduces to

$$M(t) = \frac{\alpha\lambda\Gamma(b)}{B(a,b)} \sum_{j=0}^\infty \frac{(-1)^j}{\Gamma(b-j)j!} \int_0^\infty e^{(t-\lambda)x} (1 - e^{-\lambda x})^{\alpha j} dx.$$

Changing to the variable $u = e^{-\lambda x}$, we obtain

$$M(t) = \frac{\alpha\Gamma(b)}{B(a,b)} \sum_{j=0}^\infty \frac{(-1)^j}{\Gamma(b-j)j!} \int_0^1 u^{-t/\lambda} (1-u)^{\alpha(a+j)-1} du.$$

The above expression shows that the mgf of the BGE distribution exists if $t < \lambda$. Assuming that $t < \lambda$, we have

$$M(t) = \frac{\alpha\Gamma(b)}{B(a,b)} \sum_{j=0}^\infty \frac{(-1)^j}{\Gamma(b-j)j!} B(1 - t/\lambda, \alpha(a+j)). \tag{15}$$

Analogously, if $b > 0$ is integer, assuming $t < \lambda$, we use the binomial expansion in (14) to obtain

$$M(t) = \frac{\alpha}{B(a,b)} \sum_{j=0}^{b-1} \binom{b-1}{j} (-1)^j B(1 - t/\lambda, \alpha(a+j)). \tag{16}$$

If we take $a = b = 1$ in (16), the above mgf reduces to

$$M(t) = \alpha B(1 - t/\lambda, \alpha),$$

which agrees with Gupta and Kundu's [5] equation (2.3).

From (15) and (16) with $\alpha = 1$ we have the mgf of the BE distribution

$$\begin{aligned} M(t) &= \frac{\Gamma(b)}{B(a,b)} \sum_{j=0}^{\infty} \frac{(-1)^j}{\Gamma(b-j)j!} B(1 - t/\lambda, a + j) \\ &= \frac{B(b - t/\lambda, a)}{B(a,b)}, \end{aligned} \tag{17}$$

which agrees with Nadarajah and Kotz's [11] equation (3.1). The last equality is given with more details in the Appendix A. The expressions for the mgf of the DGE distribution are obtained with $a = 1$ in (15) and (16) for $b > 0$ real non-integer and $b > 0$ integer, respectively.

We can easily obtain expressions for the mgf of the order statistics from (12) and (13). The mgf of $X_{i:n}$ for $b > 0$ real non-integer is

$$M_{X_{i:n}}(t) = \sum_{k=0}^{n-i} \sum_{m_1=0}^{\infty} \cdots \sum_{m_{k+i-1}=0}^{\infty} \delta_{k,i}^{(1)} M_{k,i}(t)$$

and for $b > 0$ integer

$$M_{X_{i:n}}(t) = \sum_{k=0}^{n-i} \sum_{m_1=0}^{b-1} \cdots \sum_{m_{k+i-1}=0}^{b-1} \delta_{k,i}^{(2)} M_{k,i}(t),$$

where $M_{k,i}(t)$ is the mgf of the $BGE(\alpha\{a(i+1) + \sum_{j=1}^{k+i-1} m_j\}, b, \lambda, \alpha)$ distribution.

4 Moments

The rth moment of the BGE distribution can be obtained from $E(X^r) = d^r M_X(t)/dt^r|_{t=0}$. Hence, if $b > 0$ is real non-integer, we have from (15)

$$\mu'_r = E(X^r) = \frac{\alpha\Gamma(b)}{\lambda^r B(a,b)} \sum_{j=0}^{\infty} \frac{(-1)^{j+r}}{\Gamma(b-j)j!} \left. \frac{d^r B(p, \alpha(a+j))}{dp^r} \right|_{p=1} \tag{18}$$

and if $b > 0$ is integer, we obtain from (16)

$$\mu'_r = E(X^r) = \frac{\alpha\Gamma(b)}{\lambda^r B(a,b)} \sum_{j=0}^{b-1} \binom{b-1}{j} (-1)^{j+r} \left. \frac{d^r B(p, \alpha(a+j))}{dp^r} \right|_{p=1}. \tag{19}$$

Equations (18) and (19) are the main results of this section and generalize the moments of the GE and BE distributions derived by Gupta and Kundu [5] and Nadarajah and

Kotz [11], respectively. The first four moments of the BGE distribution for $b > 0$ real non-integer are

$$\mu'_1 = \frac{\Gamma(a+b)}{\lambda\Gamma(a)} \sum_{j=0}^{\infty} \frac{(-1)^j (a+j)^{-1}}{\Gamma(b-j)j!} c_j, \quad \mu'_2 = \frac{\Gamma(a+b)}{\lambda^2\Gamma(a)} \sum_{j=0}^{\infty} \frac{(-1)^j (a+j)^{-1}}{\Gamma(b-j)j!} d_j,$$
$$\mu'_3 = \frac{\Gamma(a+b)}{\lambda^3\Gamma(a)} \sum_{j=0}^{\infty} \frac{(-1)^{j+1} (a+j)^{-1}}{\Gamma(b-j)j!} e_j, \quad \mu'_4 = \frac{\Gamma(a+b)}{\lambda^4\Gamma(a)} \sum_{j=0}^{\infty} \frac{(-1)^j (a+j)^{-1}}{\Gamma(b-j)j!} f_j,$$

and for $b > 0$ integer are

$$\mu'_1 = \frac{\Gamma(a+b)}{\lambda\Gamma(a)} \sum_{j=0}^{b-1} \binom{b-1}{j} \frac{(-1)^j}{a+j} c_j, \quad \mu'_2 = \frac{\Gamma(a+b)}{\lambda^2\Gamma(a)} \sum_{j=0}^{b-1} \binom{b-1}{j} \frac{(-1)^j}{a+j} d_j,$$
$$\mu'_3 = \frac{\Gamma(a+b)}{\lambda^3\Gamma(a)} \sum_{j=0}^{b-1} \binom{b-1}{j} \frac{(-1)^{j+1}}{a+j} e_j, \quad \mu'_4 = \frac{\Gamma(a+b)}{\lambda^4\Gamma(a)} \sum_{j=0}^{b-1} \binom{b-1}{j} \frac{(-1)^j}{a+j} f_j,$$

where the quantities c_j, d_j, e_j and f_j are given in the Appendix B.

Graphical representation of skewness and kurtosis for some choices of parameter b as function of parameter a, and for some choices of parameter a as function of parameter b, for fixed $\lambda = 1$ and $\alpha = 1$, are given in Figures 3 and 4, respectively. These plots show that the skewness and kurtosis curves increase (decrease) with a (b) for fixed b (a).

The rth moment of $X_{i:n}$ for $b > 0$ real non-integer is

$$E(X^r_{i:n}) = \sum_{k=0}^{n-i} \sum_{m_1=0}^{\infty} \cdots \sum_{m_{k+i-1}=0}^{\infty} \delta^{(1)}_{k,i} E(X^r_{k,i})$$

and for $b > 0$ integer

$$E(X^r_{i:n}) = \sum_{k=0}^{n-i} \sum_{m_1=0}^{b-1} \cdots \sum_{m_{k+i-1}=0}^{b-1} \delta^{(2)}_{k,i} E(X^r_{k,i}),$$

where the moments $E(X^r_{k,i})$ come from the general expansions (18) and (19) for the moments of the BGE distribution with parameters $\alpha\{a(i+1) + \sum_{j=1}^{k+i-1} m_j\}, b, \lambda$ and α.

The Shannon entropy of a random variable X is a measure of the uncertainty and is defined by $E\{-\log f(X)\}$, where $f(x)$ is the fdp of the X. For a random variable X with a BGE distribution, we obtain

$$\begin{aligned} E\{-\log f(X)\} &= -\log(\alpha\lambda) + \log B(a,b) + \lambda\mu'_1 + (1/\alpha - a)\{\Psi(a) - \Psi(a+b)\} \\ &\quad -(b-1)\{\Psi(b) - \Psi(a+b)\}, \end{aligned}$$

where μ'_1 is the mean given before.



Figure 3: Skewness of the BGE distribution as function of a for some values of b and as function of b for some values of a.



Figure 4: Kurtosis of the BGE distribution as function of a for some values of b and as function of b for some values of a.

5 Estimation and Inference

We assume that Y follows the BGE distribution and let $\theta = (a, b, \lambda, \alpha)^T$ be the parameter vector. The log-likelihood for a single observation y of Y is

$$\begin{aligned}\ell = \ell(a, b, \lambda, \alpha) &= \log\alpha + \log\lambda - \log B(a, b) - \lambda y + (\alpha a - 1)\log(1 - e^{-\lambda y}) \\ &\quad + (b - 1)\log\{1 - (1 - e^{-\lambda y})^{\alpha}\}, \quad y > 0.\end{aligned}$$

The components of the unit score vector $U = (\frac{\partial\ell}{\partial a}, \frac{\partial\ell}{\partial b}, \frac{\partial\ell}{\partial\lambda}, \frac{\partial\ell}{\partial\alpha})^T$ are

$$\begin{aligned}\frac{\partial\ell}{\partial a} &= \alpha\log(1 - e^{-\lambda y}) - \psi(a) + \psi(a + b), \\ \frac{\partial\ell}{\partial b} &= \log\{1 - (1 - e^{-\lambda y})^{\alpha}\} - \psi(b) + \psi(a + b), \\ \frac{\partial\ell}{\partial\lambda} &= \frac{1}{\lambda} - y + \frac{(\alpha a - 1)ye^{-\lambda y}}{1 - e^{-\lambda y}} - \frac{\alpha(b - 1)e^{-\lambda y}(1 - e^{-\lambda y})^{\alpha-1}y}{1 - (1 - e^{-\lambda y})^{\alpha}}, \\ \frac{\partial\ell}{\partial\alpha} &= \frac{1}{\alpha} + a\log(1 - e^{-\lambda y}) - \frac{(b - 1)(1 - e^{-\lambda y})^{\alpha}\log(1 - e^{-\lambda y})}{1 - (1 - e^{-\lambda y})^{\alpha}}.\end{aligned}$$

The expected value of the score vector vanishes and then

$$\begin{aligned}E\{\log(1 - e^{-\lambda Y})\} &= \frac{\psi(a) - \psi(a + b)}{\alpha}, \\ E[\log\{1 - (1 - e^{-\lambda Y})^{\alpha}\}] &= \psi(b) - \psi(a + b), \\ E\left\{\frac{(1 - e^{-\lambda Y})^{\alpha}\log(1 - e^{-\lambda Y})}{1 - (1 - e^{-\lambda Y})^{\alpha}}\right\} &= \frac{a\{\psi(a) - \psi(a + b)\} + 1}{\alpha(b - 1)}.\end{aligned}$$

For a random sample $y = (y_1, \ldots, y_n)$ of size n from Y, the total log-likelihood is $\ell_n = \ell_n(a, b, \lambda, \alpha) = \sum_{i=1}^{n}\ell^{(i)}$, where $\ell^{(i)}$ is the log-likelihood for the ith observation ($i = 1, \ldots, n$). The total score function is $U_n = \sum_{i=1}^{n}U^{(i)}$, where $U^{(i)}$ has the form given before for $i = 1, \ldots, n$. The MLE $\hat{\theta}$ of θ is obtained numerically from the nonlinear equations $U_n = 0$. For interval estimation and tests of hypotheses on the parameters in θ we obtain the 4×4 unit information matrix

$$K = K(\theta) = \begin{bmatrix} \kappa_{a,a} & \kappa_{a,b} & \kappa_{a,\lambda} & \kappa_{a,\alpha} \\ \kappa_{a,b} & \kappa_{b,b} & \kappa_{b,\lambda} & \kappa_{b,\alpha} \\ \kappa_{a,\lambda} & \kappa_{b,\lambda} & \kappa_{\lambda,\lambda} & \kappa_{\lambda,\alpha} \\ \kappa_{a,\alpha} & \kappa_{b,\alpha} & \kappa_{\lambda,\alpha} & \kappa_{\alpha,\alpha} \end{bmatrix},$$

where the corresponding elements are given by

$$\begin{aligned}
\kappa_{a,a} &= \psi'(a)-\psi'(a+b), \quad \kappa_{a,b}=-\psi'(a+b), \quad \kappa_{a,\lambda}=\frac{\alpha}{\lambda}T_{0,1,1,1,0}\\
\kappa_{a,\alpha} &= \frac{\psi(a+b)-\psi(a)}{\alpha}, \quad \kappa_{b,b}=\psi'(b)-\psi'(a+b), \quad \kappa_{b,\lambda}=-\frac{\alpha}{\lambda}T_{1,1,1,1,0,}\\
\kappa_{\lambda,\lambda} &= \frac{1}{\lambda^2}[1+(\alpha a-1)(T_{0,2,2,2,0}+T_{0,1,1,2,0})+\alpha(b-1)\{\alpha T_{2,2,2,2,0}+\\
&\quad (\alpha-1)T_{1,2,2,2,0}-T_{1,1,1,2,0}\}], \quad \kappa_{b,\alpha}=\frac{a\{\psi(a)-\psi(a+b)\}+1}{\alpha(b-1)},\\
\kappa_{\lambda,\alpha} &= \frac{1}{\lambda}\{aT_{0,1,1,1,0}-(b-1)(T_{1,1,1,1,0}+T_{2,1,1,1,1}+T_{1,1,1,1,1})\},\\
\kappa_{\alpha,\alpha} &= \frac{1}{\alpha^2}\{1+(b-1)(T_{2,0,0,0,2}+T_{1,0,0,0,2})\}.
\end{aligned}$$

Here, we have defined the following expectation

$$T_{i,j,k,l,m}=E[(1-V)^{-i}(1-V^{1/\alpha})^j V^{i-k/\alpha}\{\log(1-V^{1/\alpha})\}^l(\log V)^m],$$

where $V \sim Beta(a,b)$ and $i,j,k,l,m \in \{0,1,2\}$. The total information matrix is then $K_n = K_n(\theta) = nK(\theta)$.

Under conditions that are fulfilled for parameters in the interior of the parameter space but not on the boundary, the asymptotic distribution of

$$\sqrt{n}(\hat{\theta}-\theta) \ \text{ is } \ N_4(0, K(\theta)^{-1}).$$

The asymptotic multivariate normal $N_4(0, K_n(\hat{\theta})^{-1})$ distribution of $\hat{\theta}$ can be used to construct approximate confidence intervals and confidence regions for the parameters and for the hazard and survival functions. The asymptotic normality is also useful for testing goodness of fit of the BGE distribution and for comparing this distribution with some of its special sub-models using one of the three well-known asymptotically equivalent test statistics - namely, the likelihood ratio (LR) statistic, Rao score (S_R) and Wald (W) statistics.

An asymptotic confidence interval with significance level γ for each parameter θ_i is given by

$$ACI(\theta_i, 100(1-\gamma)\%) = (\hat{\theta}_i - z_{\gamma/2}\sqrt{\kappa^{\theta_i,\theta_i}}, \hat{\theta}_i + z_{\gamma/2}\sqrt{\kappa^{\theta_i,\theta_i}}),$$

where $\kappa^{\theta_i,\theta_i}$ is the ith diagonal element of $K_n(\theta)^{-1}$ for $i=1,2,3,4$ and $z_{\gamma/2}$ is the quantile $1-\gamma/2$ of the standard normal distribution.

Further, we can compute the maximum values of the unrestricted and restricted log-likelihoods to construct the LR statistics for testing some sub-models of the BGE distribution. For example, we may use the LR statistic to check if the fit using the BGE distribution is statistically "superior" to a fit using the GE distribution for a given data set. In any case, considering the partition $\theta = (\theta_1^T, \theta_2^T)^T$, tests of hypotheses of the type $H_0 : \theta_1 = \theta_1^{(0)}$ versus $H_A : \theta_1 \neq \theta_1^{(0)}$ can be performed by using any of the above three asymptotically statistics. The LR statistic for testing the null hypothesis H_0 is $w = 2\{\ell(\hat{\theta}) - \ell(\tilde{\theta})\}$, where $\hat{\theta}$ and $\tilde{\theta}$ are the MLEs of θ under H_A and H_0, respectively. Under the null hypothesis, $w \xrightarrow{d} \chi^2_q$, where q is the dimension of the vector θ_1 of interest. The LR test rejects H_0 if $w > \xi_\gamma$, where ξ_γ denotes the upper $100\gamma\%$ point of the χ^2_q distribution.

6 Application

In this section we fit BGE model to one real data set. The data set is obtained from Smith and Naylor [14]. The data are the strengths of 1.5 cm glass fibres, measured at the National Physical Laboratory, England. Unfortunately, the units of measurement are not given in the paper. The data set is: 0.55, 0.93, 1.25, 1.36, 1.49, 1.52, 1.58, 1.61, 1.64, 1.68, 1.73, 1.81, 2 ,0.74, 1.04, 1.27, 1.39, 1.49, 1.53, 1.59, 1.61, 1.66, 1.68, 1.76, 1.82, 2.01, 0.77, 1.11, 1.28, 1.42, 1.5, 1.54, 1.6, 1.62, 1.66, 1.69, 1.76, 1.84, 2.24, 0.81, 1.13, 1.29, 1.48, 1.5, 1.55, 1.61, 1.62, 1.66, 1.7, 1.77, 1.84, 0.84, 1.24, 1.3, 1.48, 1.51, 1.55, 1.61, 1.63, 1.67, 1.7, 1.78, 1.89.

The MLEs of the parameters and the maximized log-likelihood for the BGE distribution are

$$\hat{a} = 0.4125, \quad \hat{b} = 93.4655, \quad \hat{\lambda} = 0.92271, \quad \hat{\alpha} = 22.6124, \quad \hat{\ell}_{BGE} = -15.5995,$$

whereas for the BE distribution

$$\hat{a} = 17.7786, \quad \hat{b} = 22.7222, \quad \hat{\lambda} = 0.3898, \quad \hat{\ell}_{BE} = -24.1270$$

and for the GE distribution

$$\hat{\lambda} = 2.6105, \quad \hat{\alpha} = 31.3032, \quad \hat{\ell}_{GE} = -31.3834.$$

The LR statistics to test the hypotheses $H_0 : BE \times H_A : BGE$ and $H_0 : GE \times H_A : BGE$ are 17.0550 (p-value=3.63×10^{-5}) and 31.5678 (p-value=1.39×10^{-7}), respectively. Therefore, we reject the null hypothesis in both cases in favor of the BGE distribution at the significance level of 5%. The plots of the estimated densities of the BGE, BE and GE distributions fitted to the data set given in Figure 5 show that the BGE distribution gives a better fit than the other two sub-models.

7 Conclusions

We proposed the beta generalized exponential (BGE) distribution which generalizes the beta exponential distribution discussed by Nadarajah and Kotz [11] and the generalized exponential (also named exponentiated exponential) distribution introduced by Gupta and Kundu [4]. We provide a mathematical treatment of this distribution including the densities of the order statistics. We derive the moment generating function and provide infinite sums for the moments of the new distribution and of its order statistics. We discuss maximum likelihood estimation and obtain the information matrix. One application of the BGE distribution are given to show that this distribution could give better fit than other sub-models discussed in the literature. We hope this generalization may attract wider applications in reliability and biology.

References

[1] Cordeiro, G. M., Simas, A. B. and Stosic, B., 2008. The Beta Weibull distribution. *Submitted.*



Figure 5: Estimated densities of the BGE, BE and GE distributions for the data set from [14].

[2] Eugene, N., Lee, C. and Famoye, F., 2002. Beta-normal distribution and its applications. *Commun. Statist. - Theory and Methods*, 31, 497-512.

[3] Gradshteyn, I. S. and Ryzhik, I. M., 2000. Table of integrals, series, and products. *Academic Press: San Diego.*

[4] Gupta, R. D. and Kundu, D., 1999. Generalized exponential distributions. *Austral. and New Zealand J. Statist.*, 41 (2), 173-188.

[5] Gupta, R. D. and Kundu, D., 2001. Exponentiated Exponential Family: An Alternative to Gamma and Weibull Distributions. *Biometrical Journal*, 43, 117-130.

[6] Gupta, A. K. and Nadarajah, S., 2004. On the moments of the beta normal distribution. *Commun. Statist. - Theory and Methods*, 33, 1-13.

[7] Mudholkar, G. S. and Srivastava, D. K., 1993. Exponentiated Weibull family for analysing bathtub failure data. *IEEE Trans. Reliability*, 42, 299-302.

[8] Mudholkar, G. S., Srivastava, D. K. and Freimer, M., 1995. The exponentiated Weibull family. *Technometrics*, 37, 436-45.

[9] Mudholkar, G. S. and Hutson, A. D., 1996. The exponentiated Weibull family: some properties and a flood data application. *Commun. Statist. - Theory and Methods*, 25, 3059-3083.

[10] Nadarajah, S. and Kotz, S., 2004. The beta Gumbel distribution. *Math. Probab. Eng.*, 10, 323-332.

[11] Nadarajah, S. and Kotz, S., 2005. The beta exponential distribution. *Reliability Engineering and System Safety*, 91, 689-697.

[12] Nadarajah, S. and Kotz, S., 2006. The exponentiated type distributions. *Acta Appl. Math.*, 92, 97-111.

[13] Nassar, M. M. and Eissa, F. H., 2003. On the exponentiated Weibull distribution. *Commun. Statist. - Theory and Methods*, 32, 1317-1336.

[14] Smith, R. L. and Naylor, J. C., 1987. A comparison of maximum likelihood and Bayesian estimators for the three-parameter Weibull distribution. *Applied Statistics*, 36, pp. 358-369.

Appendix

We now give some details of (17). It follows for $t < \lambda$ that

$$\begin{aligned} B(b - t/\lambda, a) &= B(a, b - t/\lambda) = \int_0^1 \omega^{a-1}(1-\omega)^{b-t/\lambda} d\omega \\ &= \int_0^1 \omega^{a-1}(1-\omega)^{1-t/\lambda}(1-\omega)^{b-1} d\omega. \end{aligned}$$

If $b > 0$ is real non-integer, we obtain using (9) in $(1-\omega)^{b-1}$

$$\begin{aligned} B(b - t/\lambda, a) &= \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(b)}{\Gamma(b-j)j!} \int_0^1 \omega^{a+j-1}(1-\omega)^{1-t/\lambda} d\omega \\ &= \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(b)}{\Gamma(b-j)j!} B(a+j, 1-t/\lambda) \\ &= \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(b)}{\Gamma(b-j)j!} B(1-t/\lambda, a+j). \end{aligned}$$

Similarly for $b > 0$ integer and using the binomial expansion in $(1-\omega)^{b-1}$, we have

$$B(b - t/\lambda, a) = \sum_{j=0}^{b-1} \binom{b-1}{j} B(1-t/\lambda, a+j).$$

Hence, it is easy to see that (17) holds.

Some useful quantities are defined by

$$
\begin{aligned}
c_j &= \psi(\alpha(a+j)+1) - \psi(1), \quad d_j = c_j^2 + \psi'(1) - \psi'(\alpha(a+j)+1), \\
e_j &= -c_j[c_j^2 + 3\{\psi'(1) - \psi'(\alpha(a+j)+1)\}] + \psi''(1) - \psi''(\alpha(a+j)+1), \\
f_j &= \{c_j^2 + \psi'(1) - \psi'(\alpha(a+j)+1)\}[c_j^2 + 3\{\psi'(1) - \psi(\alpha(a+j)+1)\}] + \\
&\quad 2c_j^2\{\psi'(1) - \psi'(\alpha(a+j)+1)\} - 4c_j\{\psi''(1) - \psi''(\alpha(a+j)+1)\}.
\end{aligned}
$$